

correction dose

This is the insulin dose you take for high blood glucose during the day and at bedtime.

Meal time: 1 unit per 12mg/dL over 100

Bedtime: 1 unit per 12mg/dL over 112

blood glucose	daytime	bedtime
Under 100	0	0
100-112	1	0
113-124	2	1
125-136	3	2
137-148	4	3
149-160	5	4
161-172	6	5
173-184	7	6
185-196	8	7
197-208	9	8
209-220	10	9
221-232	11	10
233-244	12	11
245-256	13	12
257-268	14	13
269-280	15	14
281-292	16	15

continued on back

Only give correction dose if it has been **more than 2-3 hours** since the last dose of fast-acting insulin.

blood glucose	daytime	bedtime
293-304	17	16
305-316	18	17
317-328	19	18
329-340	20	19
341-352	21	20
353-364	22	21
365-376	23	22
377-388	24	23
389-400	25	24
401-412	26	25
413-424	27	26
425-436	28	27
437-448	29	28
449-460	30	29
461-472	31	30
473-484	32	31
485-496	33	32
497-508	34	33
509-520	35	34
521-532	36	35
533-544	37	36
545-556	38	37
557-568	39	38
569-580	40	39
581-592	41	40
Over 592	42	41

childrensdayton.org/diabetesresources

