

correction dose

This is the insulin dose you take for high blood glucose during the day and at bedtime.

Meal time: 1 unit per 12mg/dL over 125

Bedtime: 1 unit per 12mg/dL over 137

blood glucose	daytime	bedtime
Under 125	0	0
125-137	1	0
138-149	2	1
150-161	3	2
162-173	4	3
174-185	5	4
186-197	6	5
198-209	7	6
210-221	8	7
222-233	9	8
234-245	10	9
246-257	11	10
258-269	12	11
270-281	13	12
282-293	14	13
294-305	15	14
306-317	16	15
<i>continued on back</i>		

Only give correction dose if it has been **more than 2-3 hours** since the last dose of fast-acting insulin.

blood glucose	daytime	bedtime
318-329	17	16
330-341	18	17
342-353	19	18
354-365	20	19
366-377	21	20
378-389	22	21
390-401	23	22
402-413	24	23
414-425	25	24
426-437	26	25
438-449	27	26
450-461	28	27
462-473	29	28
474-485	30	29
486-497	31	30
498-509	32	31
510-521	33	32
522-533	34	33
534-545	35	34
546-557	36	35
558-569	37	36
570-581	38	37
581-593	39	38
Over 593	40	39

childrensdayton.org/diabetesresources

